

# Melbourn Village College

## First Aid Policy Addendum – COVID-19

<b>Last review date</b>	April 2020
<b>Responsible Officer:</b>	Principal

Any first aiders who are unable to maintain social distancing measure of 2m when attending a first aid incident should follow Government guidelines on usage of PPE:

When on duty they should ensure that they have immediate access to the minimum PPE – face mask, disposable gloves and plastic apron.

This is in addition to the normal First Aid equipment outlined in the School's First Aid Policy.

First Aiders should make themselves familiar with the guidance below in order to protect both themselves and others they are in contact with.

<https://www.gov.uk/government/publications/novel-coronavirus-2019-ncov-interim-guidance-for-first-responders/interim-guidance-for-first-responders-and-others-in-close-contact-with-symptomatic-people-with-potential-2019-ncov>

**The relevant sections are below:**

### **7. What to do if you are required to come into close contact with someone as part of your first responder duties**

#### **7.1 Personal protective equipment (PPE)**

Where it is not possible to maintain a 2 metre or more distance away from an individual, disposable gloves and a disposable plastic apron are recommended. Disposable gloves should be worn if physical contact is likely to be made with potentially contaminated areas or items.

The use of a fluid repellent surgical face mask is recommended and additional use of disposable eye protection (such as face visor or goggles) should be risk assessed when there is an anticipated risk of contamination with splashes, droplets of blood or body fluids.

When using a fluid repellent surgical face mask, you should mould the metal strap of the mask over the bridge of the nose and make sure the mask fits snugly under the chin, around or across any facial hair if present.

Clean your hands thoroughly with soap and water or alcohol sanitiser before putting on and after taking off PPE. In all circumstances where some form of PPE is used, the safe removal of the PPE is a critical consideration to avoid self-contamination. Guidance on putting on and taking off PPE is available. Use and dispose of all PPE according to the instructions and training provided by your employer or organisation.

#### **7.2 Cardiopulmonary resuscitation**

If you are required to perform cardiopulmonary resuscitation (CPR), you should conduct a risk assessment (in the Police this would be a “dynamic risk assessment”) and adopt appropriate precautions for infection control.

In adults, it is recommended that you do not perform rescue breaths or mouth-to-mouth ventilation; perform chest compressions only. Compression-only CPR may be as effective as combined ventilation and compression in the first few minutes after non-asphyxia arrest (cardiac arrest not due to lack of oxygen).

Cardiac arrest in children is more likely to be caused by a respiratory problem (asphyxia arrest), therefore chest compressions alone are unlikely to be effective.

If a decision is made to perform mouth-to-mouth ventilation in asphyxia arrest, use a resuscitation face shield where available.

Should you have given mouth-to-mouth ventilation there are no additional actions to be taken other than to monitor yourself for symptoms of possible COVID-19 over the following 14 days. Should you develop such symptoms you should follow the advice on what to do on the NHS website.

### **7.3 Providing assistance to unwell individuals**

If you need to provide assistance to an individual who is symptomatic and may have COVID-19, wherever possible, place the person in a place away from others. If there is no physically separate room, ask others who are not involved in providing assistance to stay at least 2 metres away from the individual. If barriers or screens are available, these may be used.

### **7.4 Cleaning the area where assistance was provided**

Cleaning will depend on where assistance was provided. It should follow the advice for cleaning in non-healthcare settings. Public areas where a symptomatic individual has passed through and spent minimal time in (such as corridors) but which are not visibly contaminated with body fluids can be cleaned in the usual way. However, all surfaces that a symptomatic individual has come into contact with must be cleaned and disinfected.

### **7.5 If there has been a blood or body-fluid spill**

Keep people away from the area. Use a spill-kit if available, using the PPE in the kit or PPE provided by your employer/organisation and following the instructions provided with the spill-kit. If no spill-kit is available, place paper towels/roll onto the spill, and seek further advice from emergency services when they arrive.